



Australian Government

Initial Assessment and Referral (IAR) Guidance for Mental Health

Assessment Domains and Rating Guide

Part C – Adolescents (aged 12-17)

May 2024

Initial Assessment and Referral (IAR) Guidance for Mental Health Assessment Domains and Rating Guide PART C - Adolescents (aged 12–17)

Copyright

© 2024 Commonwealth of Australia as represented by the Department of Health and Aged Care

This work is copyright. You may copy, print, download, display and reproduce the whole or part of this work in unaltered form for your own personal use or, if you are part of an organisation, for internal use within your organisation, but only if you or your organisation:

- (a) do not use the copy or reproduction for any commercial purpose; and
- (b) retain this copyright notice and all disclaimer notices as part of that copy or reproduction.

Apart from rights as permitted by the Copyright Act 1968 (Cth) or allowed by this copyright notice, all other rights are reserved, including (but not limited to) all commercial rights.

Requests and inquiries concerning reproduction and other rights to use are to be sent to the Communication Branch, Department of Health and Aged Care, GPO Box 9848, Canberra ACT 2601, or via e-mail to copyright@health.gov.au.

Permitted Uses

You may download, display, print, and reproduce the whole or part of this publication in unaltered form for:

- your own personal use;
- use within your organisation; or
- distribution and sharing with third parties.

but only if:

- you or your organisation do not use or reproduce the publication for any commercial purpose; and
- if reproduced, this copyright notice and all disclaimer notices are included as part of any reproduction.

Enquiries regarding use of this publication should be addressed to MH.IARProject@health.gov.au

This licence does not cover, and there is no permission given for, use of the Commonwealth Coat of Arms or any logos and trademarks (including the logo of the Department of Health).

Disclaimer

Use of this publication is not a substitute for professional knowledge and clinical judgement. Systems and processes for initial assessment and referral should consider the unique and personal circumstances of the individual client, including other health or social issues, their preferences and choices, and any risk or safety issues.

Use of this publication is at your own risk, and you must make your own assessment of its appropriateness for the environment and circumstances in which it is to be applied. The Department of Health assumes no legal liability or responsibility to anyone for the consequences of using or relying on the information included or incorporated in this publication.

This publication contains links to external websites that the Department of Health has no direct control over. It is the responsibility of users to make their own decisions about the accuracy, currency, reliability, and completeness of information contained on linked websites.

The contents of this publication may be updated from time to time and users are encouraged to regularly visit [IAR-DST](#) to check for updated versions of this publication.

Note on intended users

The Initial Assessment and Referral (IAR) Guidance for Mental Health and IAR Decision Support Tool (IAR-DST) is designed for use by Australian health professionals when a person presents to primary care for assistance for their mental health, or when the health professional providing the service identifies the person may be experiencing possible mental health symptoms and/or psychological distress. The IAR is intended to assist Australian health professionals to decide the most appropriate level of care a patient will need across five levels of care in an [Australian stepped care model](#).

The IAR Guidance documentation provides key information for Australian health professionals who undertake assessment and referral of people presenting to primary care who may have a mental health need.

It also provides information for mental health policy, program and service designers, and mental health and primary care service providers considering embedding the IAR into their systems, processes, and services.

Widespread adoption of the IAR 8 domain assessment framework and level of care model in the Australian healthcare system will aid in consistency of assessment and communication of patient information between practitioners, services, and systems.

Terms of use

As a condition of Your use of the Online Decision Support Tool and its documentation and guidance material (IAR Guidance), You must agree to these [Terms of Use](#) each time you use the Online Decision Support Tool and IAR Guidance. The use of this Part is subject to the Terms of Use.

About the IAR Guidance documentation and IAR-DST

The Initial Assessment and Referral for Mental Health (IAR) comprises the online IAR Decision Support Tool (IAR-DST) and Guidance documentation.

The IAR Guidance documentation includes a suite of documents providing information about the IAR and how to use it appropriately and effectively with people of different ages who present to the Australian primary care system with mental health symptoms and/or psychological distress.

The *Initial Assessment and Referral Guidance for Mental Health – PART A – General Guidance*, provides information on the IAR relevant to all age groups. It articulates the principles guiding the use of the IAR 8 domain assessment framework and IAR-DST to determine or confirm the most appropriate level of mental health treatment/care a patient requires, based on their current symptoms and circumstances. It provides information about the IAR five levels of care in the primary mental health care system and the types of services and supports associated with each level of care. It also provides general information about the online IAR-DST available at <https://iar-dst.online/#/>.

- Detailed information on the 8 domains and how to rate each domain for children, adolescents, adults and older adults is contained in the IAR Assessment Domains and Rating Guide for each age group, available as follows:
 - Part B - Children (aged 5-11)
 - Part C - Adolescents (aged 12-17) (*this document*)
 - Part D – Adults (aged 18-64)
 - Part E – Older Adults (aged 65 and over)
- Whilst the IAR Guidance documentation uses age to indicate the overall appropriateness of each rating guide, the final decision about the most appropriate rating guide to use with each person is based on the clinical judgment of the user, considering contextual and developmental factors.

A note on language

Mental health systems and services include many diverse terms and phrases that refer to people, roles and processes. Preferred terminology in mental health may vary from both the terms in policy documents and the terms other people prefer. Not all terms used have commonly agreed definitions, and not all readers will identify with the use of labels in the same way as they are presented here. A glossary of terms is provided in the *Initial Assessment and Referral Guidance for Mental Health – Part A – General Guidance*.

Contents

Note on intended users	3
Terms of use.....	3
About the IAR Guidance documentation and IAR-DST	3
A note on language	4
Contents.....	5
Table of figures.....	5
Table of practice points	5
Assessment Domains and Rating Guide – Adolescents (aged 12-17).....	7
General instructions for rating the domains	7
Overarching rules	7
The initial assessment domains – Adolescents	7
Rating guide for the initial assessment domains – Adolescents.....	10
Guide to rating each domain	10
Domain 1 – Symptom severity and distress	10
Domain 2 – Harm.....	12
Domain 3 – Functioning.....	15
Domain 4 – Impact of co-existing conditions.....	16
Domain 5 – Service use and response history.....	18
Domain 6 – Social and environmental stressors	18
Domain 7 – Family and other supports	20
Domain 8 – Engagement and motivation	21
Levels of care and using the IAR-DST	24
More information.....	24

Table of figures

Figure 1: the IAR domains	8
Figure 2: The primary and contextual initial assessment domains – Adolescents.....	9

Table of practice points

Practice point – eating disorders	12
Practice point – evaluating harm associated with suicidal thoughts, impulses, or behaviours	13
Practice point – safety planning.....	14
Practice point – assessment of adolescents with a suicide attempt history but no current suicidal indicators.....	14
Practice point – mandatory reporting.....	15
Practice point – definitions of cognitive impairment, intellectual disability, developmental delay, neurological condition, and learning and communication disorders.....	17

Practice point – childhood experiences of trauma.....	19
Practice point – bullying (online and in-person).....	20
Practice point – checking in when engagement or motivation is low	23

Assessment Domains and Rating Guide – Adolescents (aged 12-17)

The IAR Assessment and Referral Guidelines for Mental Health, Assessment Domains and Rating Guide Part C – Adolescents (aged 12-17) and IAR-DST for adolescents assists general practitioners and other clinicians to recommend the most appropriate level of care for an adolescent seeking or requiring mental health support (or when the health professional providing the service identifies the person may be experiencing possible mental health symptoms and/or psychological distress). In addition to the adolescent rating guide, the following rating guides are also available for use:

- Part B - Assessment Domains and Rating Guide – Children (aged 5-11)
- Part D - Assessment Domains and Rating Guide – Adults (aged 18-64)
- Part E - Assessment Domains and Rating Guide – Older Adults (aged 65 and over)
- Whilst the IAR Guidance uses age to indicate the overall appropriateness of each tool, the final decision about the most appropriate rating guide to use with each patient is based on the clinical judgment of the health professional conducting the assessment, taking into account contextual, developmental (such as how the individual is functioning on a social, physical, intellectual, cultural and emotional level and whether this is at, below or higher than what is typical for their chronological age) and other considerations.

General instructions for rating the domains

- The initial assessment is undertaken across 8 domains that describe clinical severity using a 5-point scale ranging from 0 to 4. Higher ratings indicate increased severity of the problem and the need for higher (more intensive) levels of care.
- Each rating within each domain is defined by one or more descriptors designated by alpha characters (a, b, c, etc.). Only one of these descriptors needs to be met for a rating to be selected.

Overarching rules

- If there is uncertainty in the ratings for the primary assessment domains that impact on the level of care appropriate for the person, the IAR user may need to pause the IAR process and seek additional information that will allow rating of the domains with confidence.
- Where uncertainty remains about ratings for the primary assessment domains even after the additional information is obtained, the person and family (where appropriate) should be supported to access an appropriate clinician or service for a more comprehensive assessment.
- IAR does not indicate the urgency of the response a person might require. Users must still consider the urgency of the response required and activate urgent assessment and care pathways if needed (as per their service model and local system policies and procedures). Users of the IAR should be familiar with local urgent assessment and care pathways.
- Unless otherwise stated, IAR users consider what has been 'typical' for the person over the past 30 days except where the person has experienced more recent or sudden changes or deterioration. Where this is the case, users should base their ratings on the more recent changes.
- The IAR should not be used as a screening tool because it cannot be used without some form of personalised assessment.

The initial assessment domains – Adolescents

The initial assessment process recommended in the IAR Guidance identifies 8 domains that should be explored in assessment and considered when determining the next steps in a referral process for

an adolescent who presents to primary care with a mental health need. These domains can be seen at figure 1.

Figure 1: the IAR domains




Figure 2 below provides a summary of the primary and contextual initial assessment domains for adolescents. Initial assessment should consider the person's current situation on all 8 domains. Each domain looks at specific factors relevant to making decisions about a level of care that is most likely suitable for the person's mental health treatment needs. The selection of the domains, and factors covered in each domain, aims to capture a limited number of key areas that a clinician would consider when determining the most appropriate services for an adolescent needing referral for mental healthcare.

Figure 2: The primary and contextual initial assessment domains – Adolescents

DOMAIN 1 Symptom severity and distress	<ul style="list-style-type: none"> • Current and past symptoms and duration. • Level of distress associated with the mental health issue. • Experience of a mental health condition. • Are symptoms improving/worsening, is distress improving/worsening, and are new symptoms emerging?
DOMAIN 2 Harm	<ul style="list-style-type: none"> • Suicidality – current and past suicidal ideation and attempts. • Intentional, non-suicidal self-harm – current and past. • Impulsive, dangerous, or risky behaviours with the potential for harm to self or others (including risks associated with the use of alcohol and other drugs). • The harm caused by abuse, exploitation, or neglect by others. • Unintentional harm to self or others arising from severe symptoms or self-neglect.
DOMAIN 3 Functioning	<ul style="list-style-type: none"> • The adolescent's ability to fulfil usual roles/responsibilities appropriate to their age, developmental level, and cultural background. • The adolescent's functioning within the family or home environment, in educational or vocational settings, with friends and peers and in the community. • The adolescent's ability to undertake basic activities of daily living appropriate to their age and developmental level (e.g., self-care, mobility, toileting, nutrition, and personal hygiene).
DOMAIN 4 Impact of co-existing conditions	<ul style="list-style-type: none"> • Physical health conditions. • Cognitive impairment, intellectual disability, developmental delay, neurological conditions, or learning and communication disorders. • Substance use.
DOMAIN 5 Service use and response history	<ul style="list-style-type: none"> • Whether the adolescent/family has previously sought help from or referred to mental health services and related supports (including specialist or mental health inpatient services). • Their progress or benefit from past or current services and support.
DOMAIN 6 Social and environmental stressors	<ul style="list-style-type: none"> • The degree to which any or all the following factors are relevant to the adolescent's current circumstances and the referral decision: significant transitions, peer group stress, trauma or victimisation, family or household stress, socio-economic disadvantage, performance-related pressure, and legal issues.
DOMAIN 7 Family and other supports	<ul style="list-style-type: none"> • Whether personal supports, including emotionally nurturing relationships, practical support, and social support, are present in the adolescent's environment and their potential to contribute to improved mental health.
DOMAIN 8 Engagement and motivation	<ul style="list-style-type: none"> • The adolescent and their parent/caregiver's awareness of the mental health issue and their capacity and willingness to engage in or accept assistance.

Rating guide for the initial assessment domains – Adolescents

The rating guide includes a hierarchical ranking of factors relevant to each domain to guide judgements about problem severity.

The rating guide provides a rating system that grades each domain on a 5-point rating scale of severity - while the terms vary in some domains, the rating scale for each domain follows the general format where:

0 = No problem

1 = Mild problem

2 = Moderate problem

3 = Severe problem

4 = Very severe problem

The rating guide outlines specific criteria for assessing each domain, designed to serve as a checklist of factors to consider when judging the extent to which a problem is present.

Guide to rating each domain

- If more than one descriptor applies to the person being assessed within each domain, the descriptor with the highest rating should be selected.
 - Example one: if 3-b and 3-c apply, but 4-a is also present, the rating selected is 4.
 - Example two: if 2-a and 2-b apply, but 3-c is also present, the rating selected is 3.
- Use all available information in making a rating. This should include clinical interviews and information gathered from the person, the person's family, referrers, or other informants where possible. Consider all reliable perspectives when selecting a rating (e.g., including information provided by the person, family, or referrer).
- The coding of ratings as numerals does not imply that an overall composite score can be used for making decisions about the person's service needs. The numbers should be regarded as simply shorthand for summarising severity.
- Guidance is given for each domain on examples of problems that should be considered for specific ratings (the 'descriptors'). Consider these as examples only rather than an exhaustive list of all factors relevant to the domain. Therefore, referring to the underlying rating format at times may be helpful.

Domain 1 – Symptom severity and distress

This domain considers symptoms to include both internalised (emotional) problems experienced by the adolescent (e.g., anxiety and depressive symptoms) as well as externalised behaviours observable by or impacting on others (e.g., impulsive behaviours, perceived concerning or aggressive behaviours, appearing to ignore instructions from adults, seeming distracted or unable to concentrate).

Symptoms may be associated with distress, but this is not always the case and (for example) may present as somatic symptoms like headaches and stomach pain. Symptoms may indicate a particular diagnostic condition, but a diagnosis is not required for rating an individual on this domain, determining an appropriate level of care, or referring the person for mental health services.

Assessment of an adolescent on this domain should consider:

- Current and past symptoms and duration.
- Level of distress associated with the mental health issues.
- Previous experience of a mental health condition.

- Are symptoms improving/worsening, is distress improving/worsening, and are new symptoms emerging?

0 = No problem in this domain

1 = Mild

Symptoms are likely to be sub-diagnostic and have been experienced for less than 3 months (but this may vary)

- Mild anxiety-related symptoms (e.g., occasional fears, worry, difficulty concentrating, body image issues, occasional unexplained somatic symptoms like headache and stomach pain) without significant avoidant behaviour.
- Mild mood-related symptoms (e.g., sadness, fatigue, apathy, some reluctance to participate in previously enjoyed activities, irritability, occasional disrupted sleep).
- Mild behavioural symptoms (e.g., distractibility, overactivity, occasional difficulty completing tasks, quick to anger, occasional concerning or aggressive behaviours, occasionally appearing oppositional, minor interpersonal difficulties).
- Currently experiencing a mental health condition associated with mild distress or mild reduction in quality of life.

2 = Moderate

Symptoms are at a level that would likely meet diagnostic criteria and have been experienced for more than 3 months (but this may vary)

- Moderate anxiety-related symptoms (e.g., excessive worry, agitation, panic, difficulty concentrating, significant self-consciousness or significant concerns about body image, appearance or weight, frequent unexplained somatic complaints) with significant avoidance of anxiety provoking situations.
- Moderate mood-related symptoms (e.g., excessive sadness, apathy, exhaustion, frequent irritability, loss of interest and pleasure and/or frequent reluctance to participate in previously enjoyed activities, frequent sleep disturbance).
- Moderate behavioural symptoms (e.g., frequent impulsivity, hyperactivity, non-adherence to age-appropriate rules or social norms, frequent concerning or aggressive behaviours, significant interpersonal difficulties).
- Currently experiencing a mental health condition associated with moderate levels of distress and/or moderate reduction in quality of life.
- History of a diagnosed mental health condition earlier in childhood that has not responded to treatment, with continuing symptoms but only associated with mild to moderate levels of distress.

3 = Severe

- Severe anxiety-related symptoms are present most of the time, the adolescent has difficulty controlling or managing the symptoms and seeks to avoid anxiety provoking situations and/or experiences severe distress if asked to engage in anxiety provoking situations such that there is severe distress and/or significant disruption to the adolescent's (and/or parent/family's) life.
- Severe mood-related symptoms are present most of the time, the adolescent has difficulty controlling or managing the symptoms and the symptoms are associated with severe distress and/or significant disruption to the adolescent's (and/or parent/family's) life.
- Significant behavioural symptoms are present most of the time the adolescent has difficulty controlling or managing the symptoms and the symptoms are associated with severe disruption and/or distress for the adolescent, and/or their parent/family and interpersonal relationships.
- Currently experiencing other severe mental health symptoms or severe psychological distress (e.g., complex trauma responses, obsessions, compulsions, severely disordered eating). Symptoms may be ongoing or of more recent or sudden onset.
- Symptoms suggestive of an early form of a severe mental health condition (e.g., odd thinking/behaviour/speech, abnormal perceptions, suspicious thinking, rapid mood swings, a substantial decrease in the need for sleep) or symptoms suggestive of an eating disorder.

- f. Has been treated by a specialist community mental health service or admitted to hospital for a mental health condition in the previous 12-months.

4 = Very severe

- a. Very severe and pervasive anxiety symptoms are present virtually all the time, the adolescent can rarely control or manage the symptoms and the adolescent refuses to engage in anxiety provoking situations or activities. The symptoms are associated with severe distress, significantly reduced quality of life and/or severe disruption to nearly all aspects of the adolescent's (and/or parent/family's) life.
- b. Very severe and pervasive mood-related symptoms are present virtually all the time, and the adolescent can rarely control or manage the symptoms. The symptoms are associated with severe distress, significantly reduced quality of life and/or severe disruption to nearly all aspects of the adolescent's (and/or parent/family's) life.
- c. Extreme behavioural symptoms are present virtually all the time, and the adolescent can rarely control or manage the symptoms. The symptoms are associated with severe distress, significantly reduced quality of life and/or severe disruption to nearly all aspects of the adolescent's (and/or parent/family's) life.
- d. Currently experiencing very severe symptoms (e.g., disordered thinking, extreme mood variation, obsessions, compulsions, extreme avoidant behaviour, extreme interpersonal difficulties, extremely disordered eating with associated physical symptoms). Symptoms may be ongoing or of more recent or sudden onset.
- e. Highly unusual and bizarre symptoms/behaviours indicating a severe mental illness (e.g., hallucinations, delusions). Symptoms may be ongoing or of more recent or sudden onset.

Practice point – eating disorders

Eating disorders are mental illnesses accompanied by physical and mental health complications which may be severe and life-threatening. A person with an eating disorder can experience disturbances in behaviours, thoughts, and feelings towards body weight/shape or food and eating. A person with symptoms suggestive of an eating disorder requires a comprehensive eating disorders assessment and referral to appropriate services according to a person's needs.

IAR-DST users should be familiar with their local eating disorder screening and assessment pathway. Contact the GP or state/territory mental health services for care instructions in the absence of a defined local pathway. For more information about eating disorders, visit the National Eating Disorder Collaboration - <https://nedc.com.au/eating-disorders/>.

Domain 2 – Harm

This domain is focused on:

- Suicidality – current and past suicidal ideation, intent, planning, and attempts.
- Intentional, non-suicidal self-harm – current and past.
- Impulsive, dangerous, or risky behaviours with the potential for psychological or physical harm to self or others (consider and include risks associated with the use of alcohol and other drugs).
- The psychological or physical harm caused by abuse, exploitation, or neglect by others.
- Unintentional harm to self, arising from symptoms or self-neglect.

*The IAR for adolescents includes the **harm from others** in Domain 2 because there are direct implications for the intensity of mental health response an adolescent at risk of or experiencing harm from others is likely to require. Placing harm from others in another domain (e.g., Domain 6) does not carry the same weight within the logic that underpins the recommendations about a level of care. Note that the presence of external stressors (e.g., family violence) is rated at Domain 6, but the degree of harm arising from those stressors is rated separately at Domain 2.*

0 = No concerns about harm

1 = Previous but no current concerns about harm

- a. No current suicidal ideation, but the adolescent has experienced suicidal ideation, plans, or intent in the past. Demonstrates future-oriented thinking and has strong protective factors.
- b. Occasional non-suicidal self-injurious acts in the recent past and not requiring any medical treatment.
- c. May have engaged in past behaviours that posed a risk to self or others, but no current or recent instances.
- d. Currently at low risk of harm from abuse, exploitation, or neglect by others.

2 = Some current concerns about harm

- a. Current suicidal ideation, without plan or intent but may have had plans, intent, or suicide attempts in the past. Demonstrates future-orientated thinking and has strong protective factors or previous suicide attempt (longer than 12 months ago) but no current ideation, intent, or plan.
- b. Frequent non-suicidal self-injurious acts in the recent past that did not require any medical treatment.
- c. Current or recent behaviours that pose a non-life-threatening risk to self or others.
- d. Currently at some risk of harm from abuse, exploitation, or neglect by others.
- e. Intermittent lapses in self-care that may lead to harm.

3 = Significant current concerns about harm

- a. Current suicidal ideation with a plan, but no current intent or a strong reluctance to carry out a plan. May have a history of suicide attempts. Strong protective factors and a commitment to engage in a safety plan, including the involvement of family, significant others, or services.
- b. Recent suicide attempt (within past 12 months) but no current ideation, intent, or plan.
- c. Frequent non-suicidal self-injurious acts in the recent past and requiring medical treatment.
- d. Recent or current impulsive, dangerous, or risky behaviours that pose a risk of harm to self or others, or that have had or are likely to have a serious negative impact.
- e. Serious medical risks and/or complications associated with a mental illness.
- f. Significant risk of, or recent experience of, abuse, exploitation, or neglect by others.
- g. Clearly compromised self-care ability that is ongoing to the extent that indirect or unintentional harm to self is likely.

4 = Very significant current concerns about harm

- a. Current suicidal ideation with intent, typically with a plan and means to carry out the plan or history of previous suicide attempt. Few or no protective factors. Limited or no future-orientated thinking.
- b. History of life-threatening self-injurious acts that are prominent in the current presentation.
- c. There is evidence of current severe symptoms (e.g., hallucinations, avoidant behaviour, paranoia, disordered thinking, delusions, impulsivity) with behaviour that is likely to present an imminent or unpredictable danger to self or others.
- d. Extremely compromised self-care ability to the extent that there is a real and present danger of the adolescent experiencing harm related to these deficits.
- e. Life-threatening medical risks and/or complications associated with a mental illness.
- f. Other signs or indicators of imminent risk of serious harm to themselves or others.

Practice point – evaluating harm associated with suicidal thoughts, impulses, or behaviours

This domain must be considered in the context of information gathered across the other seven domains. Information gathered across the other seven domains (e.g., severe symptoms, impulsivity, use of substances, environmental stressors, recent changes, and degree of engagement with helping resources) is especially important when evaluating harm.

The IAR-DST is not a suicide risk assessment or risk formulation tool. If an individual expresses suicidal thoughts or impulses or displays suicidal behaviours, a risk formulation compatible with local or state-based protocols (e.g., Towards Zero, Connecting with People) is indicated.

A risk formulation generally involves:

Determining risk status through consideration of static factors such as a history of psychiatric illness, family history of suicide, history of abuse, and history of suicidal behaviour.

Exploring risk state through consideration of recent suicidal behaviours, current symptoms and stressors, and engagement with helping resources. Comparing the current risk state to the person's "baseline" and "worst-point" states. Exploring the risk state includes building an understanding of the:

- Nature of the suicidal thoughts (frequency, intensity, speed of onset, persistence, intrusiveness)
- Perception of the future (hope, alternatives to suicide)
- Degree of planning
- Degree of preparation
- Ability to resist thoughts of suicide

Considering the **resources available** to the person and **foreseeable changes** that might exacerbate risk, a **suicide risk formulation may need to happen urgently**. If this is the case, refer to localised urgent assessment and care pathways.

Practice point – safety planning

If indicated, a safety plan can be an important resource to develop with a patient. There are templates and guidance for developing a safety plan available online from mental health service providers and systems.

Practice point – assessment of adolescents with a suicide attempt history but no current suicidal indicators

While any history of a suicide attempt is a risk factor for a subsequent attempt, several factors elevate this risk. These factors include:

1. When the previous suicide attempt occurred: the risk of re-attempting suicide remains high throughout a person's life but is particularly high in the 12-months following an attempt.
2. The lethality of the previous attempt: the method, medical seriousness, rescuability, and physical consequences of the attempt all contribute to assessing potential lethality. A previous attempt with higher lethality is associated with an increased risk of a subsequent attempt.
3. Post-attempt aftercare services: a person's participation in high-quality post-attempt aftercare services is associated with a decreased risk of a subsequent suicide attempt.

The factors contributing to a person attempting or dying by suicide are complex and highly variable. A previous suicide attempt is considered a risk factor – however, some people who attempt suicide may never again experience suicidal thinking or behaviours.

An adolescent who has previously attempted suicide is rated at minimum as a 2 (moderate risk) on this domain - or higher if there are current suicidal behaviours. A rating of 2 on Domain 2 is aligned with Level 3 care and above. This rating will help to ensure a comprehensive mental health assessment is made available (a core feature of all Level 3 services). Following a comprehensive mental health assessment, the clinician can consider the suitability of lower-intensity interventions (Level 1/Level 2).

Practice point – mandatory reporting

Mandatory reporting laws aim to identify adolescents at risk, including abuse and neglect incidents, and protect the individual adolescents involved. The laws require selected groups of people to report suspected child abuse and neglect to government authorities. Laws exist in all Australian jurisdictions. However, the laws are not the same across all jurisdictions. Differences exist in who must report, the nature of risks and incidents that must be reported, and to whom the report is made.

It is important to note that any person is lawfully entitled to make a report if they are concerned for an adolescent's welfare, even if they are not required to do so as a mandatory reporter.

Users of the IAR-DST should be familiar with signs of abuse and neglect and their legal responsibilities regarding mandatory reporting. Visit: the Australian Institute for Family Studies for more information: <https://aifs.gov.au/cfca/publications/mandatory-reporting-child-abuse-and-neglect> or seek advice from your professional indemnity insurer or professional association.

Domain 3 – Functioning

This domain considers functional impairment associated with or exacerbated by mental health issues. While some types of illnesses and disabilities experienced by the adolescent may play a role in determining what types of support services may be required, they should not be considered in determining mental health service **intensity** within a stepped care continuum.

Assessment of an adolescent on this domain should consider the impact of the mental health issues on:

- The adolescent's ability to fulfil usual roles/responsibilities appropriate to their age, developmental level, capability, and cultural background.
- The adolescent's functioning within the family or home environment, in educational, social, or vocational settings, with friends and peers and in the community.
- The adolescent's ability to undertake basic activities of daily living appropriate to their age, capability, and developmental level (e.g., self-care, mobility, toileting, nutrition, and personal hygiene).

0 = No problem in this domain

1 = Mild impact

- a. Mildly diminished ability to function in one or more of their usual roles (e.g., at home, in educational, social, or vocational settings, with friends and peers and in the community), but without significant or adverse consequences.
- b. Mental health issues contribute to brief and transient disruptions in one or more areas of functioning.

2 = Moderate impact

- a. Moderate functional impairment in more than one of their usual roles (e.g., at home, in educational or vocational settings, with friends and peers and in the community) to the extent that they are reasonably frequently unable to meet the requirements of those roles but without significant or adverse consequences.
- b. Mental health issues contribute to occasional difficulties with basic activities of daily living (e.g., eating, mobility, bathing, getting dressed, and toileting) or instrumental activities of daily living (e.g., preparing food, cleaning, transportation, managing money) but without threat to health.

3 = Severe impact

- a. Significant difficulties with functioning, resulting in disruption to many areas of the adolescent's life most of the time (e.g., limited participation in educational or vocational activities, deterioration in or some withdrawal from the community or relationships with friends and peers), but the adolescent can function independently with adequate treatment, family, and community support.
- b. Mental health issues frequently contribute to difficulties with basic activities of daily living (e.g., eating, mobility, bathing, getting dressed, and toileting) or instrumental activities of daily living (e.g., preparing food, cleaning, transportation, managing money) on a consistent basis but without threat to health.

4 = Very severe impact

- a. Profound difficulties with functioning, resulting in significant disruption to virtually all areas of the adolescent's life (e.g., unable to participate in educational, social, or vocational activities, complete withdrawal from community, friends, and peers).
- b. Mental health issues contribute to severe and persistent self-neglect that poses a threat to health.

Domain 4 – Impact of co-existing conditions

Increasingly, individuals are experiencing and managing multi-morbidity (coexistence of multiple conditions, including chronic disease). This domain considers the extent to which other conditions contribute to (or have the potential to contribute to) increased severity of the mental health issue or compromise the adolescent's ability to participate in the recommended services and support.

Assessment of an adolescent on this domain should consider the presence, and impact of, three possible coexisting conditions:

- Physical health conditions (consider all physical health issues).
- Cognitive impairment, intellectual disability, developmental delay, neurological conditions, or learning and communication disorders.
- Substance use.
- Where the adolescent has more than one of the coexisting conditions, consider the condition which has the most impact.

0 = No problem in this domain

1 = Minor impact

- a. Physical health condition(s) present but are stable and have no or a minimal impact on the adolescent's mental health.
- b. Cognitive impairment, intellectual disability, developmental delay, neurological condition, or learning and communication disorder present but has no or minimal impact on the adolescent's mental health.
- c. Recent episodes of substance use are limited, are not currently causing any concerns, and do not impact the adolescent's mental health.

2 = Moderate impact

- a. Physical health condition(s) present and moderately impacts the adolescent's mental health.

- b. Cognitive impairment, intellectual disability, developmental delay, neurological condition, or learning and communication disorder and moderately impacts, or has the potential to moderately impact the mental health of the adolescent.
- c. Occasional substance use impacts on, or has the potential to impact on, the adolescent's mental health.
- d. Non prescribed use of prescription medications impacts on, or has the potential to impact on, the adolescent's mental health.

3 = Severe impact

- a. Physical health condition(s) present, which requires intensive medical monitoring and severely impacts the adolescent's mental health (e.g., worsened symptoms, heightened distress).
- b. Cognitive impairment, intellectual disability, developmental delay, neurological condition, or learning and communication disorder present and severely impacts the adolescent's mental health.
- c. Frequent substance use threatens health and wellbeing or represents a barrier to mental health-related recovery.
- d. Non prescribed use of prescription medications significantly impacts the adolescent's mental health or presents a barrier to mental health-related recovery.
- e. Occasional use of high or extreme risk substances. (e.g., substances with a high risk of adverse outcomes such as injury, loss of life, criminal charges and/or use of injection drugs which have a high risk of infection of blood-borne diseases).

4 = Very severe impact

- a. One or more significant physical health conditions exist that are poorly managed or life-threatening and in the context of a concurrent mental health condition.
- b. Cognitive impairment, intellectual disability, developmental delay, neurological condition, or learning and communication disorder present and very severely impacts the adolescent's mental health.
- c. Regular and uncontrolled substance use.
- d. Regular and uncontrolled non-prescribed use of prescribed medications that has the potential to threaten health and well-being.
- e. Frequent use of high or extreme risk substances (i.e., substances with a high risk of adverse outcomes such as injury, loss of life, criminal charges and/or use of injection drugs which have a high risk of infection of blood-borne diseases).

Practice point – definitions of cognitive impairment, intellectual disability, developmental delay, neurological condition, and learning and communication disorders

The terms cognitive impairment, intellectual disability, developmental delay, neurological condition, and learning and communication disorders have no universally agreed definitions. For this Guidance, the below definitions will apply:

Cognitive impairment – A description of a person's current functioning regarding learning, communication, attention, memory, thinking and problem-solving. Cognitive impairment can be temporary, permanent, mild, moderate, or severe. Cognitive impairment can affect what the person can understand and how they relate to others and interpret the environment.

Intellectual disability – A disability characterised by significant intellectual functioning and adaptive behaviour limitations, covering many everyday social and practical skills. This disability originates before the age of 18. Genetic factors cause most intellectual disabilities. However, there are other causes of intellectual disabilities, such as brain injury or being born prematurely.

Developmental delay – A developmental delay is when an adolescent's physical, social, emotional, language or communication skill development is not at the level expected for their age and significantly affects their ability to engage in daily routines and activities.

Neurological condition – Neurological conditions affect the brain, spinal cord, and the nerves that connect them. There are more than 600 nervous system diseases (e.g., epilepsy, motor neurone disease, traumatic brain injury, multiple sclerosis).

Learning and communication disorders – Learning and communication disorders may affect how an adolescent comprehends, recalls, understands, or expresses information. These disorders are often dynamic and can improve over time. The impairment caused by these disorders might be minimal or significant and vary from person to person.

Domain 5 – Service use and response history

This domain considers the adolescent and their family's previous use of services and support focussed on mental health-related assistance. The initial assessment on this domain should consider:

- Whether the adolescent or their family has previously sought help from or required mental health services and related supports (including specialist or mental health inpatient services).
- Their progress or benefit from past services and support.

Definition of the term services and support - Relevant services and support refer to safe, developmentally, and culturally appropriate evidence-informed mental health, health or community services focussed on or relevant to the adolescent's mental health (such as a psychological service delivered by a GP or mental health professional or other behavioural services) rather than the personal supports provided by friends, family, or social networks.

Consider both the adolescent and their family's use of previous services and support but do not include those services and support relevant to, but not focused on, the adolescent's mental health.

0 = No previous service use

- a. Has not previously sought help or required a referral for mental health issues.

1 = Excellent progress from previous service use

- a. Previously accessed services for a mental health issue and experienced a significant benefit resulting in no need for additional services at that time.

2 = Moderate progress from previous service use

- a. Previously accessed services and experienced a moderate benefit and required some additional services (either ongoing or periodically) to maintain the benefit.

3 = Minor progress from previous service use

- a. Previously accessed services with only minor benefits resulting in a need for additional services or longer duration of services.

4 = Negligible progress from previous service use

- a. Previously accessed services with little or no benefit.

Domain 6 – Social and environmental stressors

This domain considers the extent and severity of a range of factors in the adolescent's environment that might contribute to the onset or continuation of the mental health issue. Significant environmental stressors and adversity can lead to increased symptom severity and compromise the capacity of the adolescent and their family to participate in or benefit from the recommended resources or services. Furthermore, understanding the complexities the adolescent is experiencing (or has experienced)

may alter the type of service offered or indicate that additional service referrals are required (e.g., a referral to a social support service).

Assessment on this domain should consider the degree to which any or all of the following factors are relevant to the adolescent's current circumstances and the referral decision:

- Significant losses (e.g., loss of friends or social connections, death of a loved one).
- Significant transitions (e.g., disruption to educational or vocational activities, parental separation/divorce, death of loved one, a romantic breakup, transitions relating to gender or sexual identity).
- Peer group stress (e.g., bullying, conflict with or isolation from peer group, loss of friendships).
- Trauma (e.g., emotional, physical, psychological, or sexual abuse, exploitation, witnessing or being a victim of violence, family and domestic violence, intimate partner violence, natural disaster, exposure to suicide in family/community/school or peer group, loss, conflict).
- Victimization (e.g., human rights abuses, discrimination, racial abuse, financial abuse, victim of crime, refugee, or asylum-seeking experiences).
- Family or household stress (e.g., household drug or alcohol abuse, a parent or family member with an illness or disability, carer stress or stress associated with a caregiver role).
- Performance-related pressure (e.g., unrealistic role expectations or responsibilities, schooling or work demands, caregiving responsibilities) and stressors related to high-performance demands in school, dance, sport, and other relevant extra-curricular activities.
- Socioeconomic disadvantage (e.g., poverty, parental unemployment, unstable or insecure housing).
- Legal issues (e.g., the juvenile justice system or family court involvement, enforced separation from family).

Evidence points to the contribution made by historical childhood adverse events to longer-term mental health development. Assessment on this domain should consider the adolescent's history but **only** record higher ratings where earlier experiences impact the current situation and require additional specific resources and services.

0 = No problem in this domain

1 = Mildly stressful environment

- a. The adolescent is experiencing (or has experienced) one or more stressors that are currently having or are likely to have only a minor impact on the adolescent's mental health.

2 = Moderately stressful environment

- a. The adolescent is experiencing (or has experienced) one or more stressors that are currently having or are likely to have a moderate impact on the adolescent's mental health.

3 = Highly stressful environment

- a. The adolescent is experiencing (or has experienced) one or more stressors that are currently having or are likely to have a significant impact on the adolescent's mental health.

4 = Extremely stressful environment

- a. The adolescent is experiencing (or has experienced) one or more stressors that are extreme, enduring, or recurring and are currently having, or are likely to have, a severe impact on the adolescent's mental health.

Practice point – childhood experiences of trauma

Adverse Childhood Experiences (ACEs) are stressful events or circumstances that people may experience throughout their childhood. They may relate to childhood physical, sexual, or emotional

abuse, physical or emotional neglect, exposure to family violence, parental substance use, parental mental illness, parental separation or divorce, or parental incarceration. A summary of the evidence and impacts by Emerging Minds reiterates that:

“Exposure to ACEs does not mean poor outcomes are inevitable. If present and reinforced in a child’s life, there are known protective factors that can build the child’s resilience and reduce the impacts of adversity. Nurturing relationships form the basis of healthy brain development, effective early learning, and a child’s capacity to positively respond and adapt to life challenges. Many adults who experienced significant adversity in their childhood have had successful lives and happy relationships – (Marie-Mitchell & Kostolansky, 2019; Traub & Boynton-Jarrett, 2017).

How a person responds to trauma is highly variable, and many individuals who have been exposed to ACEs will not require a mental health service. Immediate assignment of a level of care based on the experience of trauma alone is problematic and should be avoided. The ACEs study demonstrated a strong relationship between a person’s exposure to ACEs and their physical and mental health throughout their lives. Researchers have established a dose-response association for ACEs – for instance, four or more ACEs are associated with an increased risk of adverse impacts.

Practice point – bullying (online and in-person)

Bullying can impact the mental health of adolescents. Adolescents who experience bullying can experience feelings like shame, fear, embarrassment, anger, and worry. There is a marked increase in the risk of poor mental health outcomes, self-harm and suicidal ideation and behaviours among people who experience bullying, particularly if the experience of bullying is severe or prolonged.

Bullying, whilst common, is not a normal part of growing up, and an initial assessment with an adolescent should explore the adolescent’s experience of bullying and the impacts of these experiences.

Bullying should be considered in the context of social and environmental stressors (domain 6). The impacts of bullying on the adolescent (if present) will be captured in symptom severity and distress (domain 1), harm (domain 2), and functioning (domain 3).

When considering the level of care, consideration should be given to the informal supports that an adolescent might require outside the formal mental health system- social, school, family and community supports are generally important for adolescents experiencing bullying.

Domain 7 – Family and other supports

This domain considers whether personal supports, including emotionally nurturing relationships, practical support, and social support are present in the adolescent’s environment and their potential to contribute to improved mental health. This domain does not include or consider professional support. Personal supports include:

- Family/primary caregivers.
- Friends and peers.
- Supports within the school environment.
- Supports within the community (e.g., cultural connections, elders, spiritual leaders, sporting groups, neighbours etc.).

Personal supports may be present, but unable to provide the needed support at the time. There are a range of factors that may impact on whether personal supports are able to be provided, such as competing caring responsibilities, a lack of access to respite or other supports, financial or practical constraints, additional skill development requirements, or illness or distress in family or primary

caregivers. It is important to avoid blame or judgement of personal supports when exploring this domain.

Where appropriate, a mental health assessment and intervention for the support person (or family as a whole) should be considered.

0 = Highly supported

- a. There are family/primary caregivers and other personal supports available that are highly supportive, willing, and capable to meet the adolescent's developmental, emotional, practical, and social needs.

1 = Well supported

- a. There are a few family/primary caregivers and other personal supports available that are supportive, willing, and capable of meeting the adolescent's developmental, emotional, practical, and social needs.

2 = Limited supports

- a. There are a few family/primary caregivers available to provide support, but their willingness to provide support is variable or difficult to access, or the sources of support have insufficient resources or capabilities to meet the adolescent's developmental, emotional, practical, and social needs whenever it is needed, or the adolescent is reluctant to utilise the available supports.
- b. Other personal supports are available for the adolescent but only partially compensate for needs not met within the family.

3 = Minimal supports

- a. Very few actual or potential useful sources of support are available, willing, and capable of meeting the adolescent's developmental, emotional, practical, and social needs.
- b. There are serious limitations in the capacity or availability of supports outside the family, so that developmental, emotional, practical, or social needs are mostly unmet.

4 = No supports

- a. No useful sources of support are available, and developmental, emotional, practical, and/or social needs are mostly unmet.
- b. The adolescent has no access to other supports that could compensate for needs not met within the family.

Domain 8 – Engagement and motivation

This domain considers the adolescent or their parent/caregiver's awareness of the mental health issue and their motivation to engage in or accept assistance.

Many adolescents do not have the agency or resources required to seek and access services and support independently. Therefore, the engagement and motivation of the parent/caregiver is the primary determinant of access and uptake, and the parent/caregiver sub-scale is used. Whilst the parent/caregiver sub-scale rates the engagement and motivation of the parent/caregivers, the adolescent should be included in discussions, using language they understand, and supported to express their choices, preferences, fears, and goals about referral next steps.

The parent/caregiver sub-scale is used when the adolescent cannot exercise decision-making control of their healthcare decisions. The parent/caregiver sub-scale considers:

- Ability and capacity to support the adolescent to manage the condition.
- The parent/caregiver's motivation to assist the adolescent to access necessary support (critical if considering self-management options).

Conversely, where the adolescent can exercise decision-making control of their healthcare decisions, the adolescent's engagement and motivation take precedence (adolescent sub-scale). The adolescent sub-scale considers:

- The adolescent's motivation to participate in the recommended services and support.

Parent/Caregiver Sub-scale

Use the parent sub-scale where the adolescent cannot exercise decision-making control of their healthcare decisions.

0 = Optimal

- The parent/caregiver is motivated and capable of participating fully in the recommended services and supports.
- The parent/caregiver is capable of taking an active role in supporting the adolescent to manage the condition.

1 = Positive

- The parent/caregiver is mostly willing to accept and participate in the recommended services and support.
- The parent/caregiver can mostly take an active role in supporting the adolescent to manage the condition.

2 = Limited or mixed

- The parent/caregiver is unsure whether they will accept or participate in the recommended services and supports or has limited capacity to do so.
- There is significant divergence between the parents/caregivers in the level of engagement, motivation, or ability to participate in the recommended services and supports.

3 = Minimal

- The parent/caregiver cannot participate in the recommended services and support without considerable practical or emotional assistance.
- Despite the adolescent requiring them, the parent/caregiver has not facilitated access to services and supports in the past due to low engagement or motivation.

4 = Disengaged

- The parent/caregiver cannot support participation in services and supports or avoids potentially useful and available supports.

Adolescent Sub-scale

Use the adolescent sub-scale where the adolescent can exercise decision-making control of their healthcare decisions. In most instances, when working with a mature minor (see practice point on informed consent in Part A), the use of the adolescent sub-scale will be appropriate.

0 = Optimal

- a. The adolescent is motivated to participate in the recommended services and support.
- b. The adolescent is capable of taking an active role in managing the condition.

1 = Positive

- a. The adolescent is mostly willing to accept and participate in the recommended services and support.
- b. The adolescent can mostly take an active role in managing the condition.

2 = Limited or mixed

- a. The adolescent is hesitant to accept and participate in the recommended services and support.

3 = Minimal

- a. The adolescent is very reluctant to accept or participate in services and support.
- b. The adolescent has not participated in services and support in the past, despite requiring them, due to low levels of engagement or motivation.

4 = Disengaged

- a. The adolescent refuses to accept or participate in the recommended services and support.

Practice point – checking in when engagement or motivation is low

A follow-up check-in helps determine if the recommended information, resources, or services are being utilised and perceived as helpful. Proactively “checking in” or encouraging the adolescent and parent/caregiver to “check back” is essential when engagement or motivation is low. A plan for check-in should be made at the point of referral and documented.

The check-in should explore the following questions:

1. Is the adolescent engaging with the recommended information, resources, and services? If the adolescent is not engaging, it is essential to re-examine motivation and explore reasons for the lack of engagement.
2. Does the adolescent think that the recommended information, resources, and services are/were helpful?
3. Is there evidence of deterioration or changing risk of suicide or harm to self or others?
4. Is the adolescent experiencing new or worsening social and environmental stressors?
5. Discuss and document the next steps in collaboration with the adolescent. The next steps might include:
 - Continue existing service arrangements
 - Build in additional supports
 - Initiate a referral to a different level of care

Levels of care and using the IAR-DST

The IAR assists health professionals to determine the most appropriate level of mental healthcare for a person with mental health symptoms and/or psychological distress.

The IAR uses five different levels of care that are consistent with the application and understanding of the stepped care model in the Australian primary care system. The information gathered through the initial assessment and the ratings assigned against each IAR domain is used to assist in determining a level of care and inform a referral decision. The levels of care do not replace individualised assessment and care - instead, providing a framework to guide decision-making.

It is important to emphasise that the descriptions of care at each level are offered only to guide judgements about the recommended level of care. Each presenting person, parent/caregiver, and family will have unique requirements that must always take precedence in decision-making.

Health professions can use the online IAR Decision Support Tool available at <https://iar-dst.online/#/> to enter their ratings on each domain for a patient and obtain a suggested level of care.

More detail on the IAR Levels of Care and the online IAR-DST is provided in *PART A - Initial Assessment and Referral (IAR) Guidance for Mental Health – General Guidance*.

More information

Please contact the Department at MH.IARProject@Health.gov.au with questions or feedback about the IAR Guidance and IAR-DST.